



NUMBER 51 / MARCH 2011

# EUROPEAN RESPIRATORY *monograph*

CLINICAL HANDBOOKS FOR THE RESPIRATORY PROFESSIONAL

## Difficult-to-Treat Severe Asthma

Edited by K.F. Chung,  
E.H. Bel, S.E. Wenzel.



ERS

EUROPEAN  
RESPIRATORY  
SOCIETY

every breath counts

# Guest Editors



*K.F. Chung*

K.F. Chung is Prof. of Respiratory Medicine and Head of the Experimental Studies Unit at the National Heart and Lung Institute, Imperial College London (London, UK), and is an Honorary Consultant Physician at the Royal Brompton Hospital (London). K.F. Chung trained in internal and respiratory medicine in Oxford and London, and undertook postdoctoral training as an MRC visiting scientist at the University of California (San Francisco, CA, USA). He was awarded MD and DSc degrees in 1983 and 2001, respectively, from the University of London. K.F. Chung is a Senior Investigator of the UK National Institute for Health Research and co-leader of the IMI-funded EU/EFPIA UBIOPRED project on Systems Biology of Severe Asthma. He has received research grants from the Medical Research Council, the Wellcome Trust, US Health Effects Institute, EU Innovative Medicines Initiative, National Environmental Research Council (UK), Asthma UK and the National Institutes of Health. He currently serves on the editorial boards of several respiratory and pharmacological journals and has authored and co-authored more than 750 scientific articles, reviews and book chapters.



*E.H. Bel*

E.H. Bel is Prof. of Medicine and Head of Respiratory Medicine at the Academic Medical Centre (Amsterdam, the Netherlands). She qualified in medicine from the University of Amsterdam and received her PhD degree (*cum laude*) at Leiden University Medical Centre (the Netherlands). She trained in pulmonary medicine, serving as consultant and senior consultant at Leiden University Medical Centre. Her current research focuses on the more severe types of asthma that are difficult to treat, with a particular emphasis on the phenotyping, pathophysiological mechanisms, monitoring and therapeutic approach of this complex condition. Her research questions emanate directly from daily pulmonary practice and in her attempts to find answers she uses pulmonary physiological, histological and molecular research techniques. She has published extensively in high-profile journals and has been involved in the organisation of international courses and seminars. She is an active member of several respiratory societies including the ATS and the European Respiratory Society where she served as an Assembly Head and member of the Executive Committee until 2008. She is also on the editorial board of the *American Journal of Respiratory and Critical Care Medicine*, and reviews papers for many other respiratory journals.




*S.E. Wenzel*

S.E. Wenzel began her translational research career in the area of asthma in the 1980s, while a pulmonary fellow at the Medical College of Virginia (Richmond, VA, USA). She moved to the National Jewish Health (Denver, CO, USA) in 1987. She was one of the first investigators to perform invasive (bronchoscopic) studies of asthma and has continued performing pathophysiologic studies of asthma and severe asthma. She organised an American Thoracic Society (ATS) workshop which developed a definition for severe asthma that is now widely accepted. S.E. Wenzel is currently Prof. of Medicine and Director of the University of Pittsburgh Asthma Institute @UPMC/UPSOM. She has published over 100 peer reviewed papers and has been active in the ATS, serving as deputy Editor for the *American Journal of Respiratory and Critical Care Medicine*. She served as a member of the NHLBI Internal Board of Scientific Counsellors (2003–2007) and is currently on the Lung Cellular and Molecular Immunology study section. In 2006, she was given the National Jewish Faculty Award for Scientific Achievement. She received the Elisabeth Rich MD Award for serving as a role model of females in medicine/science and received the ATS 2010 Recognition Award for Scientific Accomplishments.

# Preface

---



Tremendous progress has been made in the treatment of patients with asthma during the past decades. Whereas 30 years ago patients suffering with acute asthma attacks were admitted to the emergency department several times a day, today this has been reduced to a couple of patients a year. Treatment with inhaled corticosteroids (ICS) in the first instance, followed by the use of combination therapy with ICS and long-acting  $\beta$ -agonists later on has greatly improved the long-term prognosis of asthma patients. Nevertheless, there are a number of patients that are not well controlled even with a high-dose combination therapy. Near fatal asthma attacks still occur and asthma mortality, although low, is still present. Difficult-to-treat asthma has been recognised as the severe form of the “other” asthma entities in the past. However, the understanding of the pathophysiology of this kind of asthma has been better understood during the last few years. Different phenotypes of difficult-to-treat asthma have been recognised and have led to new treatment strategies focused on pathophysiological concepts and individual clinical signs and symptoms of the patient. More and more insights into cell and molecular biology, genomics, epigenetics, epidemiology and clinical presentation of the asthma patients make it possible to start with a more personalised treatment strategy that orientates itself on the patients needs.

The Guest Editors of this issue have succeeded in attracting leading experts from within the field to write chapters and provide an overview from current pathophysiology, diagnostic strategies and treatments to future developments and what is perceived to be on the horizon.

I would like to congratulate the Guest Editors for this excellent Monograph, which will be of great interest to both basic scientists and clinicians, who will find it both insightful and resourceful in their daily practice.

Editor in Chief  
T. Welte

# Introduction

K.F. Chung<sup>\*</sup>, E.H. Bel<sup>#</sup> and S.E. Wenzel<sup>¶</sup>

<sup>\*</sup>National Heart and Lung Institute, Royal Brompton Hospital Biomedical Research Unit, Imperial College London, London, UK. <sup>#</sup>Dept of Pulmonology, University of Amsterdam, Amsterdam, The Netherlands. <sup>¶</sup>Pulmonary, Allergy and Critical Care Medicine, University of Pittsburgh, Pittsburgh, PA, USA.

Correspondence: K.F. Chung, National Heart and Lung Institute Imperial College, Dovehouse St, London, SW3 6LY, UK, Email: f.chung@imperial.ac.uk

Asthma has been recognised since early civilisation and has been well described by physicians over the centuries, with particular understanding of mechanisms from physicians in the 19th and 20th centuries [1]. For example, at the turn of the 20th century, William Osler (1849–1919) indicated in the “*Principles and Practice of Medicine*” that asthma was caused by muscular contraction, mucosal oedema and the presence of mucus in the lumen of the airways, laying the foundation of our current understanding of the nature of asthma [2]. However, real therapeutic breakthroughs only began to emerge in the latter part of the last century with the discovery of the bronchodilator effects of adrenaline and the anti-inflammatory effects of injected cortisone in the 1950s [3]. It then took nearly 20 years to demonstrate the efficacy of inhaled corticosteroids in chronic asthma [4], and a further 20 years for their use to be combined with long-acting  $\beta_2$ -adrenergic agonists making this one of the most effective treatments available for symptomatic asthma [5].

In parallel with the development in the modern treatment for asthma, there was growing concern for the increase in asthma prevalence and its severity, particularly in Western industrialised countries. Various surveys undertaken amongst asthmatics indicated a picture of continuing morbidity with increasing asthma fatalities. This led to the initiation of national guidelines for the management of asthma starting in Australia and Canada, shortly followed by several other countries and culminating in the Global Initiative for Asthma (GINA) guidelines [6]. It is likely that these guidelines have been instrumental in the reduction of death rates observed in several countries. In Finland a concerted governmental action against asthma has led to a significant reduction in morbidity and costs concerning asthma, as well as the impact on individual asthmatics and on society in general [7].

It appears that the battle for control of asthma is being won. However, amongst these improvements in asthma treatment and management approaches, which have certainly been beneficial to many asthmatics, physicians have always encountered difficult-to-treat asthma patients in their practice and these are emerging as one of the remaining challenges in asthma. This group constitutes only one in every 10 or 20 asthmatic patient and can be labelled as “difficult-to-treat severe asthma”. Another term is “therapy-resistant asthma”, which is very apt since these patients do not show significant beneficial responses to current therapies. The management challenge for the physician is to determine whether these patients are indeed asthmatics and whether they are receiving and taking the most appropriate treatments for their disease. Such patients often undergo different combinations and iterations of available treatments, sometimes at high doses and often suffering from the costly consequences of long-term systemic corticosteroid use, yet they continue to suffer from uncontrolled asthma.

A decade ago another milestone in asthma was set with the recognition of this problem of severe asthma by the European Respiratory Society (ERS) and the American Thoracic Society (ATS) with the publication of definition statements of “difficult-to-treat asthma” and on the way forward [8, 9]. The acceptance of a common definition has focused on the clinical description and phenotype of the



severe asthma patient. To make the definition more widely applicable, a global definition was recently published under a World Health Organization (WHO) consultation on severe asthma [10].

It is quite clear that the questions that needed to be answered in the 1990s were as follows. 1) What does the patient with severe asthma look like? 2) Why have these patients developed severe asthma? 3) How should existing treatments be used and what new treatments can be given to control severe asthma? 10 years on and these questions are still unanswered; however, there are now important insights from the information that has been obtained from research into severe asthma during this time. Perhaps the most pertinent information is how the disease is heterogeneous, and the challenge for the next 10 years would be to understand the different types of severe asthma.

It is also important to determine whether severe asthma represents an altogether different new form of asthma. If one espouses the generally accepted notion that asthma is the result of a complex gene–environment interaction then it could make sense. Certainly, the world has changed more rapidly over the past 25 years than it has in the previous 100 years, in terms of both the environment lived in and lifestyles. It is quite pertinent to ask whether environmental changes could have led to the increase in asthma prevalence and increasing severity of asthma, perhaps through epigenetic changes. Is it a result of disordered immunity to the new challenging environment of allergens, pollutants and infections? Would this drive the inflammatory response that leads to severe asthma? It is important that such questions and other related issues be asked and be answered.

The Guest Editors of this Monograph are fortunate to be part of an ongoing ATS/ERS Task Force on severe asthma, through which an eminent group of experts were asked to contribute to the Monograph. The task asked of them was to describe what we have learnt, what are the objectives for the next 10 years, and what milestones can be expected at the end of the next decade. We would like to thank all those who have contributed and we greatly appreciate the time and commitment they spent on this project. We would also like to thank the ERS Publications Office for their enthusiasm and hard work.

The burden of severe asthma is substantial and expectations for better treatments are high. We hope this Monograph represents a small step towards achieving these goals.

## Statement of interest

K.F. Chung has been reimbursed for travel expenses to international meetings and received honoraria for speaking at meetings sponsored by GlaxoSmithKline, AstraZeneca and Novartis. K.F. Chung has participated in advisory boards meetings of GlaxoSmithKline, Boehringer Ingelheim and Novartis, and received unrestricted research funding from GlaxoSmithKline and Pfizer; and has taken part in clinical trials organised by GlaxoSmithKline, Novartis and Asthmatyx. E.H. Bel has received a fee for speaking from GlaxoSmithKline, has received funds for research from GlaxoSmithKline and Novartis, and has received fees for consulting from MSD and Novartis. S.E. Wenzel's institute, the University of Pittsburgh, has received money for multicenter research studies from MedImmune, GlaxoSmithKline, Amgen, Cephalon and Aerovance. Monies received in each case by the University of Pittsburgh were <100,000 US\$. S.E. Wenzel received consulting funds from GlaxoSmithKline, Merck, Amira, Amgen all <10,000 US\$.

## References

1. McFadden ER Jr. A century of asthma. *Am J Respir Crit Care Med* 2004; 170: 215–221.
2. Osler W. The Principles and Practice of Medicine. New York, D. Appleton, 1921.
3. Barnes PJ. Drugs for asthma. *Br J Pharmacol* 2006; 147: Suppl. 1, S297–S303.
4. Medical Research Council. Controlled trial of effects of cortisone acetate in chronic asthma. *Lancet* 1956; 798: 167–172.

5. Bateman ED, Boushey HA, Bousquet J, *et al.* Can guideline-defined asthma control be achieved? The Gaining Optimal Asthma Control study. *Am J Respir Crit Care Med* 2004; 170: 836–844.
6. Bateman ED, Hurd SS, Barnes PJ, *et al.* Global strategy for asthma management and prevention: GINA executive summary. *Eur Respir J* 2008; 31: 143–178.
7. Haahtela T, Tuomisto LE, Pietinalho A, *et al.* A 10 year asthma programme in Finland: major change for the better. *Thorax* 2006; 61: 663–670.
8. Chung KF, Godard P, Adelroth E, *et al.* Difficult/therapy-resistant asthma: the need for an integrated approach to define clinical phenotypes, evaluate risk factors, understand pathophysiology and find novel therapies. ERS Task Force on Difficult/Therapy-Resistant Asthma. European Respiratory Society. *Eur Respir J* 1999; 13: 1198–1208.
9. Proceedings of the ATS workshop on refractory asthma: current understanding, recommendations, and unanswered questions. American Thoracic Society. *Am J Respir Crit Care Med* 2000; 162: 2341–2351.
10. Bousquet J, Mantzouranis E, Cruz AA, *et al.* Uniform definition of asthma severity, control, and exacerbations: document presented for the World Health Organization Consultation on Severe Asthma. *J Allergy Clin Immunol* 2010; 126: 926–938.